arXiv:1404.6010v1 [math.AC] 24 Apr 2014

STANLEY DEPTH OF MONOMIAL IDEALS

DORIN POPESCU

Abstract. Let $I \supsetneq J$ be two monomial ideals of a polynomial algebra over a field generated in degree $\geq d$, resp. $\geq d+1$. We study when the Stanley Conjecture holds for I/J using the recent result of [6] concerning the polarization.
Key words : Monomial Ideals, Depth, Stanley depth.
2010 Mathematics Subject Classification: Primary 13C15, Secondary 13F20, 13F55, 13P10.

Introduction

Let K be a field and $S = K[x_1, \ldots, x_n]$ be the polynomial K-algebra in n variables. Let $I \supsetneq J$ be two monomial ideals of S and suppose that I is generated by monomials of degrees $\geq d$ for some positive integer d. Using a multigraded isomorphism we may assume either that $J = 0$, or J is generated in degrees $\geq d+1$.

If I, J are squarefree monomial ideals then d is a lower bound of $\operatorname{depth}_S I/J$ by [3, Proposition 3.1] (see also [15, Lemma 1.1]). Proposition 2 gives a lower bound of $\operatorname{depth}_S I/J$ in terms of degrees also in the case when I, J are not squarefree using the polarization and the so called the canonical form of I/J (see [10]).

A Stanley decomposition of a multigraded S-module M is a finite family $\mathcal{D} = (S_l, u_l)_{l\in L}$ in which u_l are homogeneous elements of M and S_l are multigraded $K-$algebra retract of S for all $l \in L$ such that $S_l \cap \operatorname{Ann}_S u_l = 0$ and $M = \sum_{l\in L} S_l u_l$ as a multigraded $K-$vector space. The Stanley depth of $\mathcal{D}$, denoted by $\operatorname{sdepth}(\mathcal{D})$, is the depth of the $S-$module $\sum_{l\in L} S_l u_l$. The Stanley depth of M is defined as

$$\operatorname{sdepth}(M) := \max\{\operatorname{sdepth}(\mathcal{D}) \mid \mathcal{D} \text{ is a Stanley decomposition of } M\}.$$

Depth and Stanley depth behave in a different way for instance $\operatorname{depth}_S(M \oplus M') = \min\{\operatorname{depth}_S M, \operatorname{depth}_S M'\}$ while for sdepth it can happen $\operatorname{sdepth}_S(M \oplus M') > \min\{\operatorname{sdepth}_S M, \operatorname{sdepth}_S M'\}$ sometimes as seen in [5, Examples 14, 16] with the help of [9]. These results were obtained using the so called the Hilbert depth (see [1], [23]). The same notion is important also in other properties of depth and Stanley depth (see [21, Proposition 2.4]).

An upper bound for $\operatorname{depth}_S M$ could be given by the following conjecture.

Conjecture 1. *(Stanley* [22]*)* $\operatorname{depth}_S M \leq \operatorname{sdepth}_S M$.

It will be very nice if this conjecture holds for $M = I/J$. Recently Ichim, Katthän and Moyano-Fernández proved that Stanley's Conjecture holds for all factors I/J as above if and only if it holds for their polarizations [6, Theorem 4.3]. Thus we may restrict to the case when I, J are squarefree monomial ideals. Unfortunately, there are few results in this case in spite of the many papers appeared on this subject (see [3], [12], [13], [7], [14], [2], [15]). It is the purpose of our paper to study what

The support from grant PN-II-RU-TE-2012-3-0161 of Romanian Ministry of Education, Research and Innovation are gratefully acknowledged.

these few results say in the non squarefree case using [6, Theorem 4.3]. We use here the lower bound given by Proposition 2 (see Theorems 5, 6 and Proposition 4).

A particular case of this conjecture is the following one.

Conjecture 2. *Suppose that $I \subset S$ is minimally generated by some squarefree monomials $f_1, \ldots, f_r$ of degrees d, and a set E of squarefree monomials of degree $\geq d+1$. If $\operatorname{sdepth}_S I/J = d+1$ then $\operatorname{depth}_S I/J \leq d+1$.*

This conjecture is studied in [17], [18], [19], [11], [16] when $r \leq 4$ and some cases when $r = 5$ (see Theorems 3, 4). Proposition 3 proves Conjecture 2 also when $r = 6$ but $d = 1$ and $E = \emptyset$.

1. A lower bound of depth and Stanley depth

Let $S = K[x_1, \ldots, x_{n-1}]$ be the polynomial K-algebra over a field K and $J \subsetneq I \subset R$ two monomial ideals. Denote by $G(I)$, respectively $G(J)$, the minimal monomial system of generators of I, respectively J.

A very important result concerning the Stanley depth is given by [6, Corollary 4.4] and we recall it below.

Theorem 1. *(Ichim, Katthän, Moyano-Fernández) Let $J \subsetneq I$ be monomial ideals of S, and let $I^p \subset J^p$ be their (complete) polarizations. Then*

$$\operatorname{sdepth}_S I/J - \operatorname{depth}_S I/J = \operatorname{sdepth} I^p/J^p - \operatorname{depth} I^p/J^p.$$

For $i \in [n]$ let $e_i = \max_{u \in G(I) \cup G(J)} \deg_{x_i} u$ and set $e_{I/J} = \sum_{i \in [n], e_i > 0} (e_i - 1)$. We have $e_{I/J} = \operatorname{depth} I^p/J^p - \operatorname{depth}_S I/J$, that is $e_{I/J}$ is the number of the new variables necessary for polarization. Suppose that I is generated by some monomials $f_1, \ldots, f_r$ of degrees $d_{I/J}$ and a set of monomials E of degrees $\geq d_{I/J} + 1$. Then

Proposition 1. $\operatorname{depth}_S I/J \geq d_{I/J} - e_{I/J}$ *and* $\operatorname{sdepth}_S I/J \geq d_{I/J} - e_{I/J}$.

Proof. By [3, Proposition 3.1] (see also [15, Lemma 1.1]) we have $\operatorname{depth} I^p/J^p \geq d_{I/J}$ because I^p, J^p are squarefree monomial ideals. Note that by polarization the degrees of monomials are preserved. It follows that $\operatorname{depth}_S I/J = \operatorname{depth} I^p/J^p - e_{I/J} \geq d_{I/J} - e_{I/J}$. The inequality concerning sdepth is similar but easier since obviously the sdepth is $\geq d_{I/J}$ in the case of a factor of some squarefree monomial ideals. □

Example 1. Let $n = 3$, $d = 12$, $I = (x_1^3 x_2^4 x_3^5, x_1^{10} x_2^2)$. Note that $e_1 = 10$, $e_2 = 4$, $e_3 = 5$ and $e_I = 16$.

Remark 1. In the above example Proposition 1 gives $\operatorname{depth}_S I \geq -4$ which is obvious. This situation will be next improved considering the so called the canonical form of I given by [10].

We recall some definitions and results from [10].

Definition 1. The power x_n^r *enters in a monomial* u if $x_n^r | u$ but $x_n^{r+1} \nmid u$. We say that I/J is *of type* $(k_1, \ldots, k_s)$ *with respect to* x_n if $x_n^{k_i}$ are all the powers of x_n which enter in a monomial of $G(I) \cup G(J)$ for $i \in [s]$ and $1 \leq k_1 < \ldots < k_s$. I/J is *in the canonical form with respect to* x_n if I/J is of type $(1, \ldots, s)$ for some $s \in \mathbb{N}$ and we say that I/J is *the canonical form* if it is in the canonical form with respect to all variables $x_1, \ldots, x_n$.

Remark 2. Suppose that I/J is of type $(k_1, \ldots, k_s)$ with respect to x_n. It is easy to get the *canonical form* I'/J' of I/J *with respect to* x_n replacing $x_n^{k_i}$ by x_n^i whenever $x_n^{k_i}$ enters in a generators of $G(I) \cup G(J)$. Applying by recurrence this procedure for other variables we get the *canonical form* of I/J, that is with respect to all variables.

Theorem 2. *(A. Popescu* [10, Theorems 1, 2]*) Let I'/J' be the canonical form of I/J. Then* $\operatorname{sdepth}_S I'/J' = \operatorname{sdepth}_S I/J$ *and* $\operatorname{depth}_S I'/J' = \operatorname{depth}_S I/J$.

Definition 2. Let I'/J' be the canonical form of I/J and set $t_{I/J} = \max\{d_{I'/J'} - e_{I'/J'}, 0\}$ (we may have $d_{I'/J'} < e_{I'/J'}$ as shows Example 3). We call $t_{I/J}$ the *index of* I/J. When $J = 0$ we write t_I instead $t_{I/J}$ for simplicity. If I, J are squarefree monomial ideals then $t_{I/J} = d_{I/J}$.

Using the terminology defined above we get a better lower bound for sdepth and depth as in Proposition 1.

Proposition 2. $\operatorname{depth}_S I/J \geq t_{I/J}$ *and* $\operatorname{sdepth}_S I/J \geq t_{I/J}$.

Proof. By Theorem 2 we have $\operatorname{depth}_S I/J = \operatorname{depth}_S I'/J' \geq \max\{d_{I'/J'} - e_{I'/J'}, 0\} = t_{I/J}$. The second inequality holds similarly. □

Remark 3. This lower bound is easy to describe but it is not the best known lower bound. For example, when $J = 0$ then a better lower bound is given by $1 + \operatorname{size}(I)$ in the terminology of [8], [4]. More precisely, if $n = 3$, $d_I = 1$, $I = (x_1, x_2x_3) = (x_1, x_2) \cap (x_1, x_3)$ then $\operatorname{size}(I) = 1$ and $t_I = d_I$ since I is squarefree. Thus $1 + \operatorname{size}(I) > t_I$.

Remark 4. In Example 1 note that I is of type $(3, 10)$ with respect to x_1, of type $(2, 4)$ with respect to x_2 and of type (5) with respect to x_3. Then the canonical form of I is $I' = (x_1x_2^2x_3, x_1^2x_2)$. Note that I is generated by monomials of degrees 12 but in I' one generator has degree 4 and the other 3. Clearly, $e_{I'} = 2$, $d_{I'} = 3$ and so the index t_I of I is 1. Thus Proposition 2 says that $\operatorname{depth}_S I \geq 1$, which is also trivial but anyway better than what follows from Proposition 1 (see Remark 1).

2. Stanley depth of monomial ideals which are not necessarily squarefree

Suppose that I is minimally generated by some squarefree monomials $f_1, \ldots, f_r$ of degrees d for some $d \in \mathbb{N}$ and a set of squarefree monomials E of degree $\geq d+1$. Let B be the set of the squarefree monomials of degrees $d+1$ of $I \setminus J$.

We start recalling two results of [16] (see also [19] and [11]).

Theorem 3. *Conjecture 2 holds for $r \leq 4$.*

Theorem 4. *Conjecture 2 holds for $r = 5$ if there exists $j \notin \cup_{i \in [5]} \operatorname{supp} f_i$, $j \in [n]$ such that $(B \setminus E) \cap (x_j) \neq \emptyset$ and $E \subset (x_j)$.*

For simplicity we denote $t = t_{I/J}$, that is the index of I/J.

Theorem 5. *Let $J \subsetneq I$ be monomial ideals of S not necessarily squarefree and I'/J' the canonical form of I/J. Suppose that I' is generated by r' monomials $f_1, \ldots, f_{r'}$ of degree $d_{I'/J'}$ and a set E' of monomials of degree $\geq d_{I'/J'} + 1$. Let B' be the set of monomials of degree $d_{I'/J'} + 1$ from $I' \setminus J'$. Assume that* $\operatorname{sdepth}_S I/J = t + 1$. *Then the following statements hold:*

(1) *If* $r' \leq 4$ *then* $\operatorname{depth}_S I/J \leq t+1$,
(2) *If* $r' = 5$ *and there exists* $j \notin \cup_{i\in[5]} \operatorname{supp} f_i$, $j \in [n]$ *such that* $(B' \setminus E') \cap (x_j) \neq \emptyset$ *and* $E' \subset (x_j)$, *then* $\operatorname{depth}_S I/J \leq t+1$.

Proof. Let I'/J' be the canonical form of I/J. By Theorem 1 we have

$$\operatorname{sdepth} I'^p/J'^p = \operatorname{sdepth}_S I'/J' - \operatorname{depth} I'^p/J'^p + \operatorname{depth}_S I'/J' = d_{I'/J'} + 1.$$

Since I'^p is generated by r' squarefree monomials of degree $d_{I'/J'}$ and a set E'^p of squarefree monomials of degree $d_{I'/J'}+1$ we get by Theorem 3 that $\operatorname{depth} I'^p/J'^p \leq d_{I'/J'} + 1$. Hence $\operatorname{depth}_S I/J = \operatorname{depth}_S I'/J' \leq t+1$ by Theorem 2, that is (1) holds. The proof of (2) is the same using Theorem 4 instead Theorem 3. □

Remark 5. Let I be generated by some monomials $h_1, \ldots, h_r$ of degree d and a set of monomials E of monomials of degree $\geq d+1$. It is possible that I' is generated by $f_1, \ldots, f_{r'}$ of degrees $d_{I'/J'}$ with $r' > r$ and a set E' of monomials of degree $\geq d_{I'/J'} + 1$. For example when $n = 2$ and $I = (x_1^3x_2^4, x_1^{11}x_2)$ we have $r = 1$ and we see that $I' = (x_1x_2^2, x_1^2x_2)$ has $r' = 2$.

Example 2. Let $n = 2$, $d = 1$, $I = (x_1)$, $J = (x_1x_2^2)$. Then $e_1 = 1$, $e_2 = 2$, $e_{I/J} = 1$, $t = 0$ and $I^p/J^p = (x_1)/(x_1x_2y_2)$, where y_2 is the new variable from polarization. We have $I/J \cong x_1K[x_1] \oplus x_1x_2K[x_1]$ as graded K-vector spaces. Thus $\operatorname{sdepth}_S I/J = 1 = t+1$. By (1) of the above theorem we get $\operatorname{depth}_S I/J \leq 1$, the inequality being in fact an equality.

Theorem 6. *Let* $J \subsetneq I$ *be monomial ideals of* S *not necessarily squarefree. Assume that* $\operatorname{sdepth}_S I/J = t$. *Then* $\operatorname{depth}_S I/J = t$

The proof is similar to the proof of Theorem 5 using now [15, Theorem 4.3] instead Theorem 3.

Example 3. Let $n = 2$, $d = 1$, $I = (x_2)$, $J = (x_1^2x_2, x_1x_2^2)$. Then $e_1 = e_2 = e_{I/J} = 2$, $t = \max\{-1, 0\} = 0$ and $I^p/J^p = (x_2)/(x_1y_1x_2, x_1x_2y_2)$, where y_1, y_2 are the new variables from polarization. Since x_2 induces a nonzero element of the socle of I/J we see that $\operatorname{sdepth}_S I/J = 0$. Thus $\operatorname{sdepth}_S I/J = t = 0$. By the above theorem we get $\operatorname{depth}_S I/J = 0$.

3. Stanley depth of a factor of squarefree monomial ideals

The above theorem implies the following corollary.

Proposition 3. *Suppose that* $I \subset S$ *is minimally generated by* 6 *variables* $\{x_1, \ldots, x_6\}$ *and* $J \subsetneq I$ *a squarefree monomial ideal. If* $\operatorname{sdepth}_S I/J = 2$ *then* $\operatorname{depth}_S I/J \leq 2$.

Proof. By [17, Proposition 1.3] we see that there exists $c = x_6x_kx_q \notin J$ for $6 < k < q \leq n$. Let B be the set of all squarefree monomials from $I \setminus J$ and $\tilde{I}$ be the ideal generated by $x_1, \ldots, x_5$ and $\tilde{E} = B \setminus ((x_1, \ldots, x_5) \cup [x_6, c])$. Set $\tilde{J} = J \cap \tilde{I}$. Then for $j = 6$ we have $\tilde{E} \subset (x_j)$. In the following exact sequence

$$0 \to \tilde{I}/\tilde{J} \to I/J \to I/J + \tilde{I} \to 0$$

the last term is isomorphic with $(x_6)/(x_6) \cap (J + \tilde{I})$ and has depth ≥ 2 and sdepth 3 because it has just the interval $[x_6, c]$. Suppose that $\operatorname{sdepth}_S I/J = 2$. By [20, Lemma 2.2] we get $\operatorname{sdepth}_S \tilde{I}/\tilde{J} \leq 2$. When $\operatorname{sdepth}_S \tilde{I}/\tilde{J} = 1$ then it is enough

to apply [15, Theorem 4.3]. If $\operatorname{sdepth}_S \tilde{I}/\tilde{J} = 2$ and $(B \setminus \tilde{E}) \cap (x_j) \neq \emptyset$ then it is enough to apply Theorem 4.

Now suppose that $(B \setminus \tilde{E}) \cap (x_j) = \emptyset$, that is $B \cap (x_6) \cap (x_1, \ldots, x_5) = \emptyset$. In the following exact sequence

$$0 \to (x_6)/(x_6) \cap J \to I/J \to I/(J, x_6) \to 0$$

if the last term has sdepth ≥ 3 then the first term has sdepth ≤ 2 as above and so also depth ≤ 2. Otherwise, the last term has sdepth ≤ 2. But the last term is isomorphic with $(x_1, \ldots, x_5)/(x_1, \ldots, x_5) \cap J$ because $B \cap (x_6) \cap (x_1, \ldots, x_5) = \emptyset$. Thus in the exact sequence

$$0 \to (x_1, \ldots, x_5)/(x_1, \ldots, x_5) \cap J \to I/J \to I/(J, x_1, \ldots, x_5) \to 0$$

the first term has sdepth ≤ 2 and so its depth ≤ 2 by Theorem 4 when there exists $k > 6$ such that $B \cap (x_1, \ldots, x_5) \cap (x_k) \neq \emptyset$. Otherwise, $J \geq (x_1, \ldots, x_5)(x_6, \ldots, x_n)$ and we get

$$\operatorname{depth}_S (x_1, \ldots, x_5)/(x_1, \ldots, x_5) \cap J = \operatorname{depth}_{\tilde{S}} (x_1, \ldots, x_5)\tilde{S}/(x_1, \ldots, x_5) \cap J \cap \tilde{S} \leq 1$$

for $\tilde{S} = K[x_1, \ldots, x_5]$. Since the last term is isomorphic with $(x_6)/J \cap (x_6)$ it has depth ≥ 2 and the Depth Lemma ends the proof. □

Proposition 4. *Suppose that $I \subset S$ is minimally generated by 6 variables $\{x_1, \ldots, x_6\}$ and $J \subsetneq I$ is a monomial ideal not necessarily squarefree. Suppose that* $\operatorname{sdepth}_S I/J = t + 1$. *Then* $\operatorname{depth}_S I/J \leq t + 1$.

The proof is similar to the proof of Theorem 5 using now Proposition 3 instead Theorem 3.

Example 4. Let $n = 7$, $I = (x_1, \ldots, x_6)$, $J = (x_1^2, x_1x_2, \ldots, x_1x_5, x_1x_7)$. Then $t = 0$. The element $\hat{x}_1 \in I/J$ induced by x_1 is annihilated by all variables but x_6. It follows that $\operatorname{sdepth}_S I/J \leq 1$. Thus $\operatorname{sdepth}_S I/J \leq t + 1$ and so $\operatorname{depth}_S I/J \leq 1$ by Proposition 4. Note that $I^p/J^p = (x_1, \ldots, x_6)/(x_1y, x_1x_2, \ldots, x_1x_5, x_1x_7)$ has sdepth ≤ 2 because now the element of I^p/J^p induced by x_1 is annihilated by all variables but x_6, y.

References

[1] W. Bruns, C. Krattenthaler, J. Uliczka, *Stanley decompositions and Hilbert depth in the Koszul complex*, J. Commutative Alg., **2** (2010), 327-357.

[2] M. Cimpoeas, *The Stanley conjecture on monomial almost complete intersection ideals*, Bull. Math. Soc. Sci. Math. Roumanie, **55(103)** (2012), 35-39.

[3] J. Herzog, M. Vladoiu, X. Zheng, *How to compute the Stanley depth of a monomial ideal,* J. Algebra, **322** (2009), 3151-3169.

[4] J. Herzog, D. Popescu, M. Vladoiu, *Stanley depth and size of a monomial ideal*, Proc. Amer. Math. Soc., **140** (2012), 493-504.

[5] B. Ichim, A. Zarojanu, *An algorithm for computing the multigraded Hilbert depth of a module,* to appear in Exp. Math., arXiv:AC/1304.7215v2.

[6] B. Ichim, L. Katthän, J. J. Moyano-Fernández, *The behavior of Stanley depth under polarization*, arXiv:1401.4309.

[7] M. Ishaq, *Values and bounds of the Stanley depth*, Carpathian J. Math. *27* (2011), 217-224.

[8] G. Lyubeznik, *On the Arithmetical Rank of Monomial ideals*, J. Algebra **112**, 86-89 (1988).

[9] A. Popescu, *An algorithm to compute the Hilbert depth* , J. Symb. Comput., (2014), http://dx.doi.org/10.1016/j.jsc.2014.03.002, arXiv:AC/1307.6084, 2013.

[10] A. Popescu, *Depth and Stanley depth of the canonical form of a factor of monomial ideals*, to appear in Bull. Math. Soc. Sci. Math. Roumanie, arXiv:AC/1402.5826v3.

[11] A. Popescu, D. Popescu, *Four generated, squarefree, monomial ideals*, 2013, to appear in Proceedings of the International Conference "Experimental and Theoretical Methods in Algebra, Geometry, and Topology, June 20-24, 2013", Editors Denis Ibadula, Willem Veys, Springer-Verlag, 2014, arXiv:AC/1309.4986v3.
[12] D. Popescu, *Stanley depth of multigraded modules*, J. Algebra **312** (10) (2009) 2782-2797.
[13] D. Popescu, *An inequality between depth and Stanley depth*, Bull. Math. Soc. Sc. Math. Roumanie **52**(100), (2009), 377-382.
[14] D. Popescu, *Graph and depth of a square free monomial ideal*, Proceedings of AMS, **140** (2012), 3813-3822.
[15] D. Popescu, *Depth of factors of square free monomial ideals*, Proceedings of AMS,**142** (2014), 1965-1972.
[16] D. Popescu, *Stanley depth on five generated, squarefree, monomial ideals*, 2013, arXiv:AC/1312.0923v3.
[17] D. Popescu, A. Zarojanu, *Depth of some square free monomial ideals*, Bull. Math. Soc. Sci. Math. Roumanie, **56(104)**, 2013,117-124.
[18] D. Popescu, A. Zarojanu, *Depth of some special monomial ideals*, Bull. Math. Soc. Sci. Math. Roumanie, **56(104)**, 2013, 365-368.
[19] D. Popescu, A. Zarojanu, *Three generated, squarefree, monomial ideals*, 2013, arXiv:AC/1307.8292v5.
[20] A. Rauf, *Depth and Stanley depth of multigraded modules*, Comm. Algebra, **38** (2010),773-784.
[21] Y.H. Shen, *Lexsegment ideals of Hilbert depth 1*, (2012), arxiv:AC/1208.1822v1.
[22] R. P. Stanley, *Linear Diophantine equations and local cohomology*, Invent. Math. 68 (1982) 175-193.
[23] J. Uliczka, *Remarks on Hilbert series of graded modules over polynomial rings*, Manuscripta Math., **132** (2010), 159-168.

Dorin Popescu, Simion Stoilow Institute of Mathematics of Romanian Academy, Research unit 5, P.O.Box 1-764, Bucharest 014700, Romania

E-mail address: dorin.popescu@imar.ro